

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/238698716>

# Resistance in *Arabidopsis thaliana* Mediated by RPW8

Article

CITATIONS

4

READS

99

7 authors, including:



**Shunyuan Xiao**

University of Maryland, College Park

102 PUBLICATIONS 3,438 CITATIONS

SEE PROFILE



**Ozer Calis**

Akdeniz University

23 PUBLICATIONS 721 CITATIONS

SEE PROFILE



**Elaine Patrick**

University of East Anglia

23 PUBLICATIONS 1,670 CITATIONS

SEE PROFILE



**John G Turner**

University of East Anglia

69 PUBLICATIONS 8,891 CITATIONS

SEE PROFILE

Some of the authors of this publication are also working on these related projects:



Molecular Characterization of New Turkish Isolate Causes Diseases on Tomato Plants Mediated Tm-22 Resistance Genes [View project](#)



Development of Powdery Mildew Resistant non-Transgenic Cucumber Variety via CRISPR/Cas9 [View project](#)

using the expression  $s_k$  to convey the meaning  $m_i$ . Similarly,  $Q_{ki} = \mu(s_k, m_i) / \sum_j \mu(s_k, m_j) = \mu(m_i, s_k)$  is the probability of interpreting the expression  $s_k$  to mean  $m_i$ . The need to communicate meanings is related to events in the shared world of the linguistic community. Therefore, one can define a measure  $\sigma$  on the set of possible meanings ( $\Sigma_m$ ) that speakers and hearers might wish to communicate with each other. Given this, we can define  $a_{ij} = \text{tr}[\rho^{(i)} \Lambda (Q^{(j)})^T]$ , where  $\Lambda$  is a diagonal matrix such that  $\Lambda_{ii} = \sigma(m_i)$ . This is the probability that an event occurs and is successfully communicated from a user of  $G_i$  to a user of  $G_j$ .  $F(G_i, G_j)$  is the probability that users of  $G_i$  will have a successful communication with each other. Communication might break down in one of two ways: (i) poverty: an event happens whose meaning cannot be encoded by  $G_i$ , and (ii) ambiguity: an event happens whose meaning has an ambiguous encoding in  $G_i$  leading to a possibility of misunderstanding. Thus,  $F(G_i, G_j)$  is a number between 0 and 1 and denotes the fitness of  $G_i$ . Maximum fitness,  $F(G_i, G_i) = 1$ , is achieved by grammars that can express every possible meaning (zero poverty) and have no ambiguities.

33. To study the effect of finite (small) population sizes, the deterministic Eq. 1 is replaced by a stochastic process. In this case, we observe that the population adopts one of the candidate grammars (that admits a stable equilibrium) for some time and then jumps to another equilibrium. If the candidate grammars differ in their fitness, then the stochastic process performs an evolutionary optimization on the space of all grammars.

34. Denote by  $x_i$  the fraction of individuals who use  $G_i$  of universal grammar  $U_1$ ; denote by  $y_i$  the fraction of individuals who use  $G_i$  of universal grammar  $U_2$ .  $U_1$  and  $U_2$  contain, respectively,  $n_1$  and  $n_2$  candidate grammars. Some of the candidate grammars can be part of both universal grammars. The universal grammars,  $U_1$  and  $U_2$ , can also differ in the number of sample sentences,  $b_1$  and  $b_2$ , that are being considered. Therefore, we have to take into account the rate of producing offspring with grammatical communication; this rate is given by the declining function  $r(b)$ . An alternative interpretation is that  $r(b)$  describes the cost that is associated with learning. The dynamics are described by

$$\dot{x}_i = r(b_1) \sum_{j=1}^{n_1} x_j f_j^{(1)} Q_{ji}^{(1)} - \phi x_i \quad i = 1, \dots, n_1$$

$$\dot{y}_i = r(b_2) \sum_{j=1}^{n_2} y_j^{(2)} f_j^{(2)} Q_{ji}^{(2)} - \phi y_i \quad i = 1, \dots, n_2$$

We have  $f_i^{(m)} = \sum_{j=1}^{n_1} x_j F(G_j^{(m)}, G_i^{(1)}) + \sum_{j=1}^{n_2} y_j F(G_j^{(m)}, G_i^{(2)})$ ,  $m \in \{1, 2\}$ , and  $\phi = \sum_{i=1}^{n_1} f_i^{(1)} x_i r(b_1) + \sum_{i=1}^{n_2} f_i^{(2)} y_i r(b_2)$ , where the superscripts 1 and 2 refer to  $U_1$  and  $U_2$ , respectively.

35. In general, it is advantageous to reduce the size of the search space, because a smaller  $n$  leads to a larger accuracy of grammar acquisition. The situation is more complex, however. Consider two universal grammars  $U_1$  and  $U_2$  with  $n_1 > n_2$ . Suppose  $U_1$  is resident and  $U_2$  is an invading mutant. If  $n_1$  exceeds the coherence threshold, then  $U_2$  will always out-compete  $U_1$ . If  $n_1$  is below the coherence threshold, then  $U_2$  can only invade if the specific grammar adopted by the population of  $U_1$  speakers is also part of  $U_2$ ; otherwise  $U_1$  can resist invasion by  $U_2$ . The selective difference between  $U_1$  and  $U_2$  is small if both  $n_1$  and  $n_2$  values are either well above or well below the coherence threshold. Hence, selection is strongest close to the coherence threshold (if  $n_1 \approx n_2$ ).

36. This problem has been solved before, in a different context. How many words,  $N$ , can be stably maintained in a population if each child hears  $b$  words during its language acquisition period and has a probability,  $p$ , to memorize a new word after one encounter? The answer is  $N < bp$  [M. A. Nowak, J. B. Plotkin, V. A. A. Jansen, *Nature* **404**, 495 (2000)].

37. The implicit assumption here is, of course, that the rule-based grammars can generate at least these  $N$  sentence types. In a principles and parameters frame-

work, condition 4 implies that the number of parameters  $k$  has to be less than  $N$ .

38. W. von Humboldt, *Linguistic Variability and Intellectual Development* (Univ. of Pennsylvania Press, Philadelphia, 1972).

39. Support from the Packard Foundation, the Leon Levy

and Shelby White Initiatives Fund, the Florence Gould Foundation, the Ambrose Monell Foundation, the Alfred P. Sloan Foundation, and the NSF is gratefully acknowledged.

21 September 2000; accepted 10 November 2000

## Broad-Spectrum Mildew Resistance in *Arabidopsis thaliana* Mediated by *RPW8*

Shunyuan Xiao, Simon Ellwood, Ozer Calis, Elaine Patrick, Tianxian Li, Mark Coleman, John G. Turner\*

Plant disease resistance (*R*) genes control the recognition of specific pathogens and activate subsequent defense responses. We show that the *Arabidopsis thaliana* locus *RESISTANCE TO POWDERY MILDEW8 (RPW8)* contains two naturally polymorphic, dominant *R* genes, *RPW8.1* and *RPW8.2*, which individually control resistance to a broad range of powdery mildew pathogens. Although the predicted *RPW8.1* and *RPW8.2* proteins are different from the previously characterized *R* proteins, they induce localized, salicylic acid-dependent defenses similar to those induced by *R* genes that control specific resistance. Apparently, broad-spectrum resistance mediated by *RPW8* uses the same mechanisms as specific resistance.

The majority of characterized *R* genes participate in gene-for-gene interactions, in which the *R* product appears to act as a receptor that recognizes a product of the corresponding avirulence (*Avr*) gene from the pathogen, inducing defense responses. The *R* gene-mediated defenses typically involve a rapid, localized necrosis, or hypersensitive response (HR), at the site of infection, and the localized formation of antimicrobial chemicals and proteins that restrict growth of the pathogen (1, 2). Many crops rely on *R* genes for resistance to specific pathogens, but resistance fails in the presence of strains of the pathogen that lack the corresponding *Avr* genes. Broad-spectrum disease resistance is therefore desirable, and has been achieved through the use of recessive mutations (3); a challenge is to develop broad-spectrum resistance with dominant *R* genes (4). More than 20 of the *R* genes that confer specific resistance have been characterized and they form five classes of protein with differing combinations of five conserved structural motifs. With the exception of Hm1, a toxin reductase, and Pto, a protein kinase, the characterized *R* proteins contain a leucine-rich repeat (LRR) motif believed to specify recognition (1, 5, 6). All of the characterized *A. thaliana* *R* genes encode proteins with motifs for a nucleotide-binding site (NBS) and an LRR, and similar *R* genes have been isolated from

other plant species (7). Probes for conserved sequences in NBS-LRR motifs have detected numerous homologs in the genomes of crop plants (7), and more than 100 in the genome of *A. thaliana* (8). The *A. thaliana* loci *RPW7* and *RPW8* from accession Ms-0 map to the same interval on chromosome 3, and confer resistance to the powdery mildew pathogens *Erysiphe cruciferarum* UEA1 and *E. cichoracearum* UCSC1, respectively (9). During the mapping of *RPW8*, we discovered that this locus segregated from the mapped NBS-LRR *R*-gene homologs (8), suggesting the presence of a different type of resistance gene, which we characterize here.

We genetically mapped *RPW8* to a fragment of genomic DNA from Ms-0 in cosmid B6 (Fig. 1A) (10), and confirmed that Col-0 plants containing the B6 transgene (T-B6) were resistant to *E. cichoracearum* UCSC1 (Fig. 2A) (10). The B6 DNA sequence (11) revealed three open reading frames (ORFs) (Fig. 1A) encoding a serine/threonine protein kinase 2 (*SPK-2*) (GenBank accession number S56718) and two uncharacterized genes, which we named *MSC1* and *MSC2*. Subclones of B6 in a plant transformation vector were introduced into Col-0 plants by *Agrobacterium*-mediated transformation (11, 12), and those that contained either *MSC1* or *MSC2*, or both of these ORFs, conferred resistance to *E. cichoracearum* UCSC1 (Fig. 1A). This indicated that *RPW8* comprises two independently acting genes, *MSC1* and *MSC2*, which we therefore renamed *RPW8.1* and *RPW8.2*, respectively. Ms-0 cDNAs for *RPW8.1* and *RPW8.2* (13) were introduced

School of Biological Sciences, University of East Anglia, Norwich NR4 7TJ, UK.

\*To whom correspondence should be addressed. E-mail: j.g.turner@uea.ac.uk

## REPORTS

into accession Col-0 plants under control of the highly active cauliflower mosaic virus 35S promoter (11, 12), and transgenic plants T-35S::RPW8.1 and T-35S::RPW8.2, homozygous for the corresponding transgene, were resistant to *E. cichoracearum* UCSC1, whereas T-35S::SKP-2 plants used as controls were susceptible (Fig. 1B) (10).


The characterized *R* genes normally control resistance to only a narrow range of isolates of a species of pathogen (1). To examine the range of pathogens controlled by *RPW8*, we inoculated *A. thaliana* Col-0 plants transgenic for *RPW8* with a selection of pathogens virulent on accession Col-0. Transgenic plants T-B6, T-35s::RPW8.1, and T-35s::RPW8.2 were resistant to all of the tested powdery mildew pathogens of *Arabidopsis*. These included 15 isolates of *E. cichoracearum*, *E. cruciferarum* isolate UEA1 [indicating that *RPW7* (9) is also determined by *RPW8.1* and *RPW8.2*], *E. orontii* isolate MGH, representing three distinct species (14), and *Oidium lycopersici* isolate Oxford. A feature of these powdery mildew pathogens is that they each have a very wide host range. For example, *E. cichoracearum* UCSC1 also infects members of the Cucurbitaceae (15), *E. cruciferarum* UEA1 also infects members of the Brassicaceae (9, 15), and *O. lycopersici* Oxford attacks members of the Cucurbitaceae and the Solanaceae (16). T-B6 plants were susceptible to the other pathogens tested, including *Peronospora parasitica* Noco2 (10), the cauliflower mosaic virus, and the bacterium *Pseudomonas syringae* pv *tomato* DC3000 (17).

The genomic sequence of *RPW8.1* contained a single 197-nucleotide (nt) intron and produced a 711-nt transcript with 444 nt of predicted coding sequence, and the genomic sequence of *RPW8.2* contained a 128-nt intron and produced a 798-nt transcript with 522 nt of predicted coding sequence (GenBank accession number AF273059). Northern analysis indicated that *RPW8.1* and *RPW8.2* were expressed constitutively, but increased after infection with *E. cichoracearum* UCSC1 (10). The predicted proteins RPW8.1 and RPW8.2 had 45.2% sequence identity (10), and were relatively small (molecular weights 17,000 and 19,973, respectively) and basic (isoelectric points of 9.46 and 10.05, respectively). Their sequences predicted an NH<sub>2</sub>-terminal transmembrane (TM) domain, or possibly a signal peptide, and a coiled-coil (CC) domain (10). Although RPW8.1 and RPW8.2 had no significant homology to any characterized proteins, they had 22 and 33% identity, and 51 and 62% similarity, respectively, to a 130 amino acid sequence forming the NH<sub>2</sub>-terminus of the predicted product of resistance protein-like gene BAB08633, which has an NBS and an LRR motif at residues 171 to 751.


We compared the DNA sequences of the *RPW8.1* and *RPW8.2* alleles from seven *A. thaliana* accessions resistant and susceptible to *E. cichoracearum* UCSC1 (11, 15). Resistance of accessions Kas-1 and Wa-1 to *E. cichoracearum* UCSC1 is controlled by loci *RPW10* and *RPW13*, respectively, both of which map to the *RPW8* locus (18). The DNA sequences of the *RPW8.1* and *RPW8.2* alleles from Kas-1 and from Wa-1 were identical to

those from Ms-0. Accessions Ler, Nd-0, and Ws-0 were moderately susceptible, and the DNA sequences of their *RPW8.1* and *RPW8.2* alleles were predicted to encode proteins with 90.5 to 98.3% amino acid sequence identity to those encoded by the Ms-0 alleles. There is an insertion at the 3' end of the allele for *RPW8.1* in Nd-0, and a premature stop at the 3' end of the alleles for *RPW8.2* in Ws-0 and Sy-0. Other differences from Ms-0 were

**Fig. 1.** Isolation of the *RPW8* locus. (A) Restriction sites in cosmid B6 used for making the indicated subclones (11). Subclones introduced into Col-0 by *Agrobacterium*-mediated transformation are marked (+) if they conferred resistance to *E. cichoracearum* UCSC1, and (–) if they did not. ORFs detected in the B6 sequence (11) are shown as thick lines. Only subclones containing either or both *MSC1* and *MSC2* conferred resistance. (B) cDNAs for *RPW8.1* and *RPW8.2* (13), and for *SKP-2* were cloned in vector pKMB (26) under control of the highly active viral 35S promoter, and introduced into *A. thaliana* Col-0 by *Agrobacterium*-mediated transformation (11, 12). Plants containing the 35S::RPW8.1 or 35S::RPW8.2 transgenes were resistant (10), indicated by (+), and plants containing the 35S::SKP-2 transgene were susceptible, indicated by (–), to *E. cichoracearum* UCSC1. (C) Organization of *RPW8.1* and *RPW8.2* and homologs in accessions Ms-0 and Col-0. Exons are indicated as thick lines; each gene has a single intron indicated by a gap.



**Fig. 2.** Analysis of *RPW8*-mediated resistance of *A. thaliana* to *E. cichoracearum*. (A) Ten days after plants were inoculated with conidia from *E. cichoracearum* UCSC1 (27), (left to right) white mycelium and vegetative spore masses of the pathogen had developed on leaves of accession Col-0, which was susceptible, but not on leaves of Col-0 plants containing the *RPW8* transgene, T-B6, which was resistant. T-B6 plants in the *eds1-2/eds1-2* background (20) were susceptible, T-B6 plants in the *ndr1-1/ndr1-1* mutant background (20) were resistant, and T-B6 plants containing the *NahG* transgene encoding salicylate hydroxylase were susceptible. (B) Germinated conidia (c, blue) on surfaces of leaves 30 hours after inoculation. Hydrogen peroxide formation was detected with diamino benzidine (28) as a brown reaction product (r) in T-B6 (right) epidermal cells penetrated by the pathogen, but could not be detected in Col-0 (left) epidermal cells penetrated (p) by the pathogen. Bar, 30 μm. (C) Northern analysis indicated that defense gene *PR-1* was induced in T-B6 (B) leaves 48 hours after inoculation with *E. cichoracearum* UCSC1, but not in Col-0 (C) leaves.





conserved: for example, in RPW8.2, two (T64S and D116G) of the five amino acid differences occurred in all four susceptible accessions (10). Accession Col-0 is extremely susceptible to mildew and we could not detect alleles of either *RPW8.1* or *RPW8.2* by Southern analysis (10). However, the corresponding sequence in Col-0 [bacterial artificial chromosome (BAC) T20E23] revealed a single gene, CAB62477, predicted to encode a protein with 52 and 33% amino acid sequence identity to *RPW8.1* and *RPW8.2*, respectively.

Some *R*-gene loci consist of clusters of homologs that represent sites of rapidly evolving *R*-gene specificity (19). Analysis of the nucleotide sequence of cosmid J4-2 from Ms-0 (10) revealed that *RPW8* was linked to three ORFs with 62.6 to 82.3% DNA sequence identity to *RPW8.1* and *RPW8.2*. These three paralogs were named Homologous to *RPW8* 1 (*HR1*), *HR2*, and *HR3* (GenBank accession number AF273059). They are closely related (99.4 to 99.9% DNA sequence identity) to their orthologs CAB62474, CAB62475, and CAB62476, respectively, in BAC T20E23 from accession Col-0 (10); we named CAB62477 as *HR4* (Fig. 1C). A recombination breakpoint between *RPW8.2* and *HR3*, detected with marker 3B3-L (10) indicated that *HR1*, *HR2*, and *HR3* did not confer resistance to the powdery mildews. The *RPW8* locus of Ms-0 therefore contained five *RPW8* paralogs, and the locus in Col-0 contained four (Fig. 1C).

A majority of the characterized *A. thaliana* *R* genes mediate resistance mechanisms that involve an HR, induced H<sub>2</sub>O<sub>2</sub> formation and expression of the pathogenesis-related (PR) genes, and require salicylic acid (*I*) and the signal pathway genes *EDS1* or *NDR1* (20). We examined whether these features occur in *RPW8*-mediated resistance. Defense responses in T-B6 plants could be detected 30 hours after inoculation with *E. cichoracearum* UCSC1, as the formation of H<sub>2</sub>O<sub>2</sub> in epidermal cells penetrated by the pathogen (Fig. 2B); the penetrated cells subsequently collapsed, forming microscopic lesions characteristic of the HR and there was no further growth of the pathogen (Fig. 2A) (9). By contrast there was no evidence of cellular H<sub>2</sub>O<sub>2</sub> when Col-0 epidermal cells were penetrated by the pathogen (Fig. 2B), which grew to form masses of white mycelia and conidia on the leaf surface in 10 days (Fig. 2A). *PR-1* transcripts accumulated in T-B6 plants 48 hours after inoculation, but not in inoculated Col-0 plants (Fig. 2B). Resistance was abolished in T-B6 plants containing salicylate hydroxylase, which converts salicylic acid to catechol (Fig. 2A). Resistance was also abolished in plants containing the *RPW8* transgene from T-B6 crossed into the *eds1-2/eds1-2* mutant background, but not in the

*ndr1-1/ndr1-1* mutant background (Fig. 2A), indicating that *EDS1*, but not *NDR1*, was required for resistance.

*RPW8.1* and *RPW8.2* are similar to many of the characterized *R* genes in that they are naturally occurring, polymorphic, dominant alleles (1), occur in a gene cluster (1, 19, 21), and induce defense response associated with HR. Unlike other characterized *R* genes, *RPW8.1* and *RPW8.2* conferred resistance to a wide range of powdery mildew diseases of *A. thaliana*, reminiscent of that conferred by recessive alleles at the barley *MLO* locus (3). However, *mlo*-based resistance involves a spontaneous-lesion phenotype and cell wall apposition in epidermal tissues before pathogen attack (3), which is different from the HR phenotype triggered by *R* genes (1, 9). None of the tested powdery mildew pathogens could overcome *RPW8*-mediated resistance, indicating that *RPW8*-mediated resistance does not involve a gene-for-gene interaction (2, 21, 22), or possibly that each of these pathogens has an *Avr* gene that interacts with *RPW8*. The NH<sub>2</sub>-termini of NBS-LRR *R* proteins play a role in the regulation of cell death (23), defense pathway-signaling (20), and in the determination of specificity (24). *RPW8.1* and *RPW8.2* had limited homology to the predicted NH<sub>2</sub>-terminus of an NBS-LRR *R*-like protein. *RPW8.1* and *RPW8.2* may therefore have some NBS-LRR *R* gene functions, to recognize powdery mildew pathogens, or to recruit NBS-LRR *R* gene-mediated defenses for resistance. Alternatively *RPW8.1* and *RPW8.2* may interact with NBS-LRR *R* proteins to initiate resistance responses, possibly as targets for pathogen virulence factors that are guarded by NBS-LRR *R* proteins as has been proposed for Pto (25).

References and Notes

1. K. E. Hammond-Kosack, J. D. G. Jones, *Annu. Rev. Plant Physiol. Plant Mol. Biol.* **48**, 575 (1997).
2. N. T. Keen, *Annu. Rev. Genet.* **24**, 447 (1990).
3. R. Buschges et al., *Cell* **88**, 695 (1997).
4. C. M. Rommens, G. M. Kishore, *Curr. Opin. Biotechnol.* **11**, 120 (2000).
5. B. J. Staskawicz, F. M. Ausubel, B. J. Baker, J. G. Ellis, J. D. Jones, *Science* **268**, 661 (1995).
6. J. L. Dangl, *Cell* **80**, 363 (1995).
7. D. Leister et al., *Theor. Appl. Genet.* **98**, 509 (1999).
8. M. A. Botella et al., *Plant J.* **12**, 1197 (1997).
9. S. Xiao, S. Ellwood, K. Findlay, R. P. Oliver, J. G. Turner, *Plant J.* **12**, 757 (1997).
10. Supplementary material is available at www.sciencemag.org/cgi/content/full/291/5501/118/DC1.
11. *Arabidopsis thaliana* accession Ms-0 genomic DNA in the plant transformation vector SLJ75515 was propagated in *Escherichia coli* strain DH10B (GIBCO-BRL). Overlapping Eco RI and Hind III fragments of cosmids B6 and J4-2 were ligated into appropriate sites in pBluescript II SK<sup>+</sup> (Stratagene), cloned in *E. coli* strain XL-Blue (Stratagene), and sequenced. *RPW8* alleles from *Arabidopsis* accessions were amplified by polymerase chain reaction (PCR) from genomic DNA with primers specific for *RPW8.1* (GACCCGTACAGTACTAGTCTA and GATTTCGGAAATGTATACAAGAA) and for *RPW8.2* (AACTCTTCACCTCGAGAGCTAACA and AGTCGTTTGACACAATTGGGACAT). Products from four independent PCRs were pooled and se-

quenced. DNA sequences were assembled with the Staden DNA analysis package and analyzed with programs found at the Human Genome Mapping Project Resource Centre (<http://www.hgmp.mrc.ac.uk>). Restriction sites detected in the sequence of cosmid B6 were used to make subclones in vector SLJ75515 propagated in *E. coli* strain DH10B. *RPW8.1* and *RPW8.2* cDNAs were amplified by RT-PCR using Pfu-Turbo (Stratagene) with primers for *RPW8.1* (CCGGAATTCATGCCGATGGTGAGCTTGCGATA and CGCGGATCCTCAAGCTCTATTTTACTACAAGC) and *RPW8.2* (CCGGAATTCATGATGTGCTGAGGTTGCCGCA and CCGGATCCTCAAGAATCATCACTGCAGAACGT), and cloned into the Eco RI-Bam HI site of pKMB for expression under control of the constitutive viral 35S promoter in *Arabidopsis* Col-0. Clones were maintained in *E. coli* DH10B. *Agrobacterium tumefaciens* strain GV3101 was transformed with plasmids by electroporation, and used for stable transformation of *Arabidopsis* accession Col-0.

12. S. J. Clough, A. F. Bent, *Plant J.* **16**, 735 (1998).
13. Transcription start and stop sequences were determined by 5'-RACE and 3'-RACE according to the manufacturer's instruction (GIBCO-BRL). Gene-specific primers for *RPW8.1* were: 5'-RACE: CCACAATATTATGCTTCT and nested primer GAACCAA-AACCGGCTCGATACTAA and 3'-RACE: AATGCACATAAAGTTCGTGAAGT. Gene-specific primers for *RPW8.2* were 5'-RACE: GGATCGCACGGTTTGT and nested primer CTGAAGTCTTGGCTACGTTTCT, and 3'-RACE: GCTAATACGATGGTGGTAGAT and nested primer CGATGGGTGGTAGATGGGATGTT. PCR products were cloned into pGEM-T easy vector (Promega) cloned in *E. coli* strain XL-Blue and sequenced. The Wisconsin sequence analysis program GAP was used to calculate the percent identity and similarity of proteins.
14. G. S. Saenz, J. W. Taylor, *Can. J. Bot.* **77**, 150 (1999).
15. L. Adam et al., *Mol. Plant-Microbe Interact.* **12**, 1031 (1999).
16. J. M. Whipps, S. P. Budge, J. S. Fenlon, *Plant Pathol.* **47**, 36 (1998).
17. S. Xiao, J. G. Turner, unpublished results.
18. S. Somerville, personal communication.
19. M. Parniske et al., *Cell* **91**, 821 (1997).
20. N. Aarts et al., *Proc. Natl. Acad. Sci. U.S.A.* **95**, 10306 (1998).
21. C. M. Thomas, M. S. Dixon, M. Parniske, C. Golstein, J. D. Jones, *Philos. Trans. R. Soc. London Ser. B* **353**, 1413 (1998).
22. B. Baker, P. Zambryski, B. Staskawicz, S. P. Dinesh-Kumar, *Science* **276**, 726 (1997).
23. C. F. Hwang, A. V. Bhakta, G. M. Truesdell, W. M. Pudlo, V. M. Williamson, *Plant Cell* **12**, 1319 (2000).
24. J. E. Luck, G. J. Lawrence, P. N. Dodds, K. W. Shepherd, J. G. Ellis, *Plant Cell* **12**, 1367 (2000).
25. E. A. vanderBiezen, J. D. G. Jones, *Trends Biochem. Sci.* **23**, 454 (1998).
26. J. Mylne, J. R. Botella, *Plant Mol. Biol. Rep.* **16**, 257 (1998).
27. *E. cichoracearum* UCSC1 was propagated on squash, *E. cruciferarum* UEA1 on oilseed rape, and *O. lycopersici* Oxford and *E. orontii* MGH were propagated on *Lycopersicon esculentum* cv Monymaker.
28. H. Thordal-Christensen, Z. G. Zhang, Y. D. Wei, D. B. Collinge, *Plant J.* **11**, 1187 (1997).
29. We thank F. M. Ausubel for *E. orontii* MGH, M. Bardin for *E. cichoracearum* isolates, J. R. Botella for pKMB, S. Covey for cauliflower mosaic virus infections, X. Dong for *Arabidopsis* Col-0 transgenic for *NahG*, B. Feys and J. Parker for *P. parasitica* Noco2 and *eds1-2*, J. Jones for pSLJ75515, Ohio State University's *Arabidopsis* Stock Center for BAC filters containing IGF and TAMU libraries, R. Oliver for discussion, P. Repetti and J. Staskawicz for *ndr1-1*, K. Schrick for CAPS marker g19397, S. Somerville for *E. cichoracearum* UCSC1, M. Stammers for yeast artificial chromosome clones and BAC filters, Z. Zhang and S. Gurr for *O. lycopersici* Oxford, and the Biology and Biotechnology Research Council, UK, for financial support.

8 September 2000; accepted 29 November 2000